



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

Barred Owl, Great Horned Owl, Yellow-billed Cuckoo, Hairy Woodpecker, Whippoorwill, Nighthawk, Blue Jay, Grasshopper Sparrow, Bank Swallow, Black-and-white Warbler, Water-Thrush, Louisiana Water-Thrush and White-breasted Nuthatch.

A noteworthy feature of bird-seeing this season was the arrival in April of warblers and other birds never previously recorded for this section before May. A marked decrease in the number of many warblers is also evident, especially Magnolia, Blackburnian, Chestnut-sided and Canadian. Of the latter none has been seen by either of us, though ordinarily not rare here. The cold and severe storms of the nesting season in the Adirondacks and Canada last year appear to have prevented normal maintenance of numbers in these and probably other species.

OTTO McCREARY AND F. H. HALL.

A NOTE ON THE SPRING MIGRATION, 1901, AT LELAND, UNCOMPAHGRE UTE INDIAN RESERVATION, UTAH.

BY H. TULLSEN.

The above is a long title for a short note, but having to do with a region so little studied ornithologically, this note may not be altogether useless. Leland, Utah, is an elevated point, about 5,000 feet above sea-level, on the Uinta, a branch of Green River. Sage-brush abounds on the dry plains, and along the streams are cotton-woods and other trees.

March 2d—Magpies first seen carrying twigs for nest.

March 17th—Flock of Mountain Bluebirds. On the day before a bird was seen perched on a post in the distance; this was probably one of these birds. From the 17th, common.

March 22d—Two western Meadowlarks seen. Next day five individuals were observed, and from that time they were common.

March 31st and April 7th and 8th—The Flycatcher (*Tyrannus verticalis*), one individual each time, was seen. Cannot state when they became common.

April 1st—Kildeers seen; they became common at once.

April 7th—Notes of a Robin heard.

April 13th—A belted Kingfisher seen. The "rattle" of one was heard a few days before this—date not recorded.

April 29th—Black-headed Grosbeaks. Several noted. They became common immediately.